



# Is There A Relationship Between The Components Of Metabolic Syndrome And Depression?

Gity Sotoudeh<sup>1</sup>, Maryam Khosravi<sup>1, 2\*</sup>, Mahdieh Hosseinzadeh<sup>3</sup>, Firoozeh Raisi<sup>4</sup>, Reza Majdzadeh<sup>5</sup>

1. Department of Community Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of medical Sciences, Tehran, Iran
2. Department of Nutrition, School of Medicine, Mashhad University of medical Sciences, Mashhad, Iran; Department of Public Health, North Khorasan University of medical Sciences, Bojnurd, Iran
3. Department of Nutrition, Shahid Sadoughi University of Medical Sciences, Yazd, Iran
4. Department of Psychiatry, Roozbeh Hospital and Psychiatry and Psychology Research Centre, Tehran University of Medical Sciences, Tehran, Iran
5. Department of Statistics and Epidemiology, School of Public Health and Knowledge Utilization Research Center, Tehran University of medical Sciences, Tehran, Iran

ARTICLE INFO	ABSTRACT
<p><i>Article type:</i> Research Paper</p> <hr/> <p><i>Article History:</i> Received: 20 Aug 2018 Accepted: 09 Dec 2018 Published: 27 Jan 2019</p> <hr/> <p><i>Keywords:</i> Depression Metabolic Syndrome MetS Waist Circumference Blood Pressure Triglyceride High Density Lipoprotein Fasting Blood Glucose</p>	<p><b>Introduction:</b> Metabolic syndrome is characterized by at least three out of five main criteria, including high waist circumference, hypertension, high level of triglyceride, low level of high-density lipoprotein, and elevated fasting blood glucose. The International Diabetes Federation has estimated the prevalence of metabolic syndrome at 25% across the world. Metabolic syndrome is more prevalent in the individuals with depression, as well as other psychiatric disorders. However, the current findings in this regard are conflicting. The present study aimed to evaluate the associations between the criteria of metabolic syndrome and depression as the most common psychiatric disorder.</p> <p><b>Methods:</b> This cross-sectional study was conducted on 130 participants aged 18-65 years. Metabolic syndrome was assessed based on the criteria of the International Diabetes Federation, and depression was evaluated based on the criteria of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).</p> <p><b>Results:</b> After the adjustment of the important confounders (physical activity, calorie intake, history of depression, depression score, menopausal status, body mass index, and dietary patterns), the waist circumference was observed to be significantly higher in the patients with depression compared to the other subjects.</p> <p><b>Conclusion:</b> According to the results, visceral obesity may lead to depression and vice versa. Therefore, it is recommended that further investigation be conducted in this regard in order to confirm this association.</p>

► *Please cite this paper as:*

Sotoudeh G, Khosravi M, Hosseinzadeh M, Raisi F, Majdzadeh R. Is There A Relationship Between The Components Of Metabolic Syndrome And Depression?. *J Nutrition Fasting Health*. 2018; 6(3): 139-144. DOI: 10.22038/JNFH.2018.34334.1139

## Introduction

The prevalence of metabolic syndrome has increased from 1988 to 2012 in various socio demographic groups. By 2012, more than one-third of the adults in the United States (34.2%) corresponded with the definition and criteria of metabolic syndrome, which have been agreed upon by several international organizations. The International Diabetes Federation has reported the prevalence rate of metabolic syndrome to be 25% across the

world. The major causes of the spread of this disorder are the increased consumption of high-calorie, low-fiber fast foods and decreased physical activity (1).

Depression is considered to be the most common psychological disorder (2). The population of the patients with depression has been estimated at 322 million globally (3). As a result, the motto of the World Health Day in 2017 was "Depression: Let's talk." (4).

\* *Corresponding author:* Maryam Khosravi, Department of Nutrition, Faculty of Medicine, Pardise Daneshgah, Azadi Square, Mashhad, Iran. Tel: 00985138002367; Email: Khosravim@mums.ac.ir

© 2018 mums.ac.ir All rights reserved.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Moreover, depression in adults has been associated with high morbidity and mortality, the increased risk of various diseases (5-8), and low personal satisfaction (9).

Recently, metabolic syndrome has become more prevalent among psychiatric patients (10). Metabolic syndrome is characterized by at least three of five main criteria, including high waist circumference, hypertension, high level of triglyceride, low level of high-density lipoprotein (HDL) and cholesterol, and elevated fasting plasma glucose. These issues are correlated with cardiovascular diseases and diabetes type II (11, 12).

Previous studies have indicated that metabolic syndrome is prevalent in severely depressed patients, while other findings have denoted bilateral relation between depression and metabolic syndrome (13, 14). Due to the controversy regarding this issue (14, 15), extensive research is required to clarify the association between the criteria of metabolic syndrome and depression.

The present study aimed to assess the correlations between the criteria of metabolic syndrome and depression.

## Material and methods

### *Subjects and Study Design*

This cross-sectional study was conducted on 130 participants at Imam Hossein and Baharloo psychiatric clinics in Tehran, Iran. The participants were enrolled for the measurement of metabolic syndrome markers and diagnosis of depression. Depressed patients were diagnosed with major depressive disorder by psychiatrists based on the criteria of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) and had no history of depression within the past year.

The inclusion criteria of the study were the age of 18-65 years and residence in Tehran, Iran. The exclusion criteria were as follows: 1) diagnosis of cognitive impairment or other psychotic disorders by a psychiatrist; 2) severe depression or inability to cooperate and complete the questionnaire; 3) receiving antidepressants or treatment for depression; 4) hormonal disorders (e.g., Addison's disease, Cushing's disease, hyperthyroidism, hypothyroidism, and hyperparathyroidism); 5) chronic diseases

(e.g., cancer, heart disease, diabetes, stroke, fibromyalgia, kidney or liver failure, multiple sclerosis, and Parkinson's disease); 6) chronic infections (e.g., HIV, mononucleosis, tuberculosis, viral hepatitis, and pneumonia) within the past two weeks; 7) alcohol/drug addiction at the time of the study or withdrawal within the past three months; 8) body mass index (BMI) of  $\geq 40$  kg/m<sup>2</sup> and 9) pregnancy and lactation at the time of the study or within the past year.

After explaining the objectives of the research, written informed consent was obtained from the participants. The study protocol was approved by the Ethics Committee of Tehran University of Medical Sciences, Iran.

### *Assessment of Covariates*

A demographic questionnaire was used to collect data, as well as some confounders. After 10 minutes of relaxation, systolic and diastolic blood pressure was measured with the subjects seated after 10 minutes relaxation in chairs with no exposure to stress using a standard mercury sphygmomanometer. Waist circumference was also measured in all the participants. These covariates were considered as the criteria for metabolic syndrome. In addition, depression was diagnosed based on the 4th edition of the DSM-IV criteria (16), which have been standardized for the Iranian population (17).

Based on the metabolic equivalent task hours per day (MET-h/day), a classified physical activity questionnaire was utilized consisting of nine activity levels (from rest/sleep [MET=0.9] to vigorous physical activity [MET $\geq$ 6]). The questionnaire has been designed in Europe and was validated using the daily physical activity questionnaire and the Computer Science Application (CSA) Accelerometer (Model 7164 Ambulatory Monitor) (18). Kelishadi et al. have confirmed the validity and reliability of the questionnaire for the Iranian population (19).

In order to measure fasting blood glucose and lipid profile (i.e., metabolic syndrome criteria), blood samples (5 ml) were collected from the subjects who had been fasting for 12 hours (7-10 AM). Glucose and lipid profile, including triglyceride (TG), HDL, very low-

density lipoprotein, and total cholesterol, were assayed using enzymatic methods (Pars Azmouk, Iran) and an automatic Selecta E device (Vitalab, Netherlands). Since the highest TG concentration in our participants was lower than 400mg/dl, the concentration of low-density lipoprotein (LDL) was calculated using the Friedewald formula, as follows (20):

$$LDL\ Cholesterol\ (mg/dl) = Total\ Cholesterol - HDL\ Cholesterol - 0.2\ TG$$

Insulin was assayed with caution using the radiation survey (Diasource, Belgium) and the antibodies labeled with a gamma counter (model: Gen II, Factory Genesis, USA). Insulin sensitivity was measured based on the following formula (21):

$$QUICKI\ Index: 1 / (\log [insulin]) (\mu U/ml) + \log (glucose) (mg/dl)$$

### Statistical Analysis

The sample size was calculated for all the criteria of metabolic syndrome, and the final sample size was determined based on the largest possible sample size and practical conditions of the researcher.

Data analysis was performed in SPSS version 20 (Chicago, IL) using the Kolmogorov-Smirnov test to assess the normality of the covariates and t-test or Mann-Whitney U test to compare the covariates between the two groups based on normality. In addition, simple logistic regression was applied to investigate the correlation between depression and metabolic syndrome criteria with an estimated odds ratio (OR). The goodness of fit (GOS) for logistic regression was examined using logical confidence intervals and Hosmer-Lemeshow test.

### Results

No significant differences were observed between the study groups in terms of quantitative factors, such as age, height, weight, BMI, and frequency of weekly tobacco use and smoking. However, patients with depression had significantly lower physical activity compared to the control group ( $P < 0.01$ ). In addition, total energy intake was significantly higher in the patients with depression compared to the controls ( $P < 0.05$ ) (Table 1).

**Table 1.** Comparison of Variables between Patients with and without Depression

variables	Depressed (Mean±SE)	Non-depressed (Mean±SE)	P value
Age (year)	35.85±1.04	35.69±0.72	0.8
Height(cm)	162.8±0.79	163.4±0.59	0.5
Weight(kg)	69.7±1.3	70.1±0.96	0.8
BMI(kg/m <sup>2</sup> )	26.4±0.49	26.4±0.37	0.9
Smoking Habits (frequency per week)	8.2±2.8	8.2±2.8	0.09
Tobacco Use (frequency per week)	0.4±0.22	0.08±0.02	0.1
Physical Activity(MET-h/day)	36.9±0.52	38.6±0.33	0.007
Total Energy Intake(kcal/day)	2887±112	2634±69.9	0.04

P-value: simple logistic regression

In current research, we assessed the association between depression and the criteria of metabolic syndrome. According to the findings, none of the variables of metabolic syndrome were significant in the study groups (Table 2). However, after the adjustment of the confounders (physical activity, calorie intake, history of depression, depression score,

menopausal status, BMI, and dietary patterns), waist circumference was observed to be significantly higher in the patients with depression compared to the controls ( $P = 0.009$ ). In other words, waist circumference was associated with depression independent of the confounders.

**Table 2.** Relation between Depression and Criteria of Metabolic Syndrome before and after Adjustment of Confounders

Criteria of Metabolic Syndrome and Other Variables	Depressed (Mean±SE)	Non-depressed (Mean±SE)	P-value <sup>a</sup>	P-value <sup>b</sup>	P-value <sup>c</sup>	OR(95%CI)
Waist Circumference (cm)	89.1±1.2	88.2±0.9	0.5	-	0.009	1.49(1.12-2.007)
Systolic Blood Pressure (mmHg)	100.6±1.6	104.9±1.2	-	0.056	0.6	0.99(0.94-1.04)
Diastolic Blood Pressure (mmHg)	64±14	66±12	-	0.36	0.8	0.99(0.93-1.06)
Fasting Blood Glucose(mg/dl)	92±599	9371±12	-	0.4	0.2	2.03(0.67-6.10)
Triglyceride(mg/dl)	101±46	98±52	0.4	-	0.9	0.99(0.97-1.02)
HDLc (mg/dl)	48.5±1.4	48.7±1.2	0.7	-	0.8	1.14(0.39-3.36)

Total Cholesterol(mg/dl)	179±4.5	177.7±4.6	0.1	-	0.5	0.72(0.23-2.25)
LDLc (mg/dl)	112.5±3.8	108.7±3.7	0.5	-	0.8	1.13(0.39-3.2)
Insulin (µIU/ml)	13.2±9	13.9±9	-	0.7	0.6	0.95(0.74-1.21)
Insulin Sensitivity	0.33±0.024	0.33±0.023	0.5	-	0.3	0.22(0.01-4.7)

P<sub>value</sub><sup>a</sup>: t-test; P<sub>value</sub><sup>b</sup>: Mann-Whitney U test; P<sub>value</sub><sup>c</sup>: multiple logistic regression after adjusting confounders (physical activity, daily calorie intake, dietary patterns, history of depression, depression score, menopausal status, and BMI)

## Discussion and Conclusion

The present study aimed to assess the correlations between depression and the criteria of metabolic syndrome, including waist circumference, HDL-C, LDL-C, TG, fasting blood sugar, systolic and diastolic blood pressure, and insulin sensitivity. According to the obtained results, only waist circumference was significantly higher in the patients with depression compared to the controls, while this association was masked by the presence of the confounders before adjustment. After adjusting some important confounders (e.g., physical activity, daily calorie intake, dietary patterns, history of depression, and depression score), the difference was revealed. Therefore, it could be concluded that waist circumference is associated with depression independent of these confounders.

Consistent with our findings, the study by Ra and Kim (22) indicated that depression was significantly associated with high waist circumference among women after the adjustment of the covariates (education level, family income, high alcohol consumption, eating out, obesity, menopause, and use of antidepressants). Therefore, it could be inferred that the association of waist circumference and depression does not depend on the mentioned confounders.

In another research in this regard, Hiles et al. reported a positive correlation between depression and antidepressant use with waist circumference. In the mentioned prospective study, the association between depression and the components of metabolic syndrome was not considered significant after the discontinuation of antidepressants. Furthermore, the discontinuation of antidepressants led to the reduction of the waist circumference in the patients with depression after four years (23). This finding could be attributed to the high activity of the parasympathetic nervous system, which is associated with worsening of metabolic control (24).

In a bidirectional prospective study (23), the

severity of depression was reported to be associated with increased waist circumference. However, the criteria of metabolic syndrome did not lead to depression. This could be due to an unhealthy lifestyle, especially smoking habits and alcohol consumption, which often persist in the patients with depression despite the resolution of the other symptoms.

According to Kahl et al. (25), Dunbar et al. (26), and Pan et al. (14), metabolic syndrome is more prevalent in the patients with depression compared to normal individuals. Moreover, the results of a large, population-based study of Mexican-Americans indicated high rates of metabolic syndrome (46%) and depression (29%) (27).

In the present study, no associations were observed between depression and the other criteria of metabolic syndrome. This could be due to the fact that all the patients had only recently been diagnosed with depression and used no antidepressants. Considering that antidepressants cause metabolic disturbances, this outcome was expected. This finding is consistent with the study by Herva et al. (29), which showed no correlations between metabolic syndrome and depression after adjusting BMI and demographic indicators. Therefore, it could be concluded that waist circumference is a significantly different criterion between depressed and healthy individuals and could be proposed for the screening of the components of metabolic syndrome in the patients with depression (28).

Owing to the cross-sectional design of the current research, the obtained data could lead to the understanding of the differences in the metabolic syndrome criteria in depressed and healthy individuals. The correlations between the criteria of metabolic syndrome and depression were assessed before and after the adjustment of some confounders and risk factors of depression. The adjusted confounders were similar to the confounders in the previous studies in this regard (15, 29), with the exception of dietary patterns, which should be taken into account (30). It is also noteworthy

that the patients in the present study had only recently been diagnosed with depression and used no antidepressants. Therefore, this factor was statistically controlled rather than adjusted. Additionally, the inclusion and exclusion criteria of the study were determined with utmost precision.

One of the limitations of the current research was that we could not realize the temporal association between depression and metabolic syndrome.

### Conflict of interest

None declared.

### References

- Moore JX, Chaudhary N, Akinyemiju T. Metabolic Syndrome Prevalence by Race/Ethnicity and Sex in the United States, National Health and Nutrition Examination Survey, 1988–2012. *Prev Chronic Dis.* 2017; 14: E24.
- Sobocki P, Ekman M, Ågren H, Krakau I, Runeson B, Mårtensson B, et al. Health-related quality of life measured with EQ-5D in patients treated for depression in primary care. *Value Health.* 2007; 10(2): 153-60.
- World Health Organization. Depression and Other Common Mental Disorders: Global Health Estimates. Geneva: World Health Organization; 2017. License: CC BY-NC-SA 3.0 IGO.
- WHO. Depression: Let's talk. 7 April 2017; <http://www.who.int/campaigns/world-health-day/2017/en> (Access Date 15 August 2018).
- Mezuk B, Eaton WW, Golden SH. Depression and osteoporosis: epidemiology and potential mediating pathways. *Osteoporos Int.* 2009; 19(1): 1-12.
- Donini LM, Savina C, Cannella C. Eating habits and appetite control in the elderly: the anorexia of aging. *Int psychogeriatr.* 2003; 15(1): 73-87.
- Palsson S, Skoog I. The epidemiology of affective disorders in the elderly: a review. *Int Clin psychopharmacol.* 1997; 12(Suppl): S3-13.
- Penninx BW, Leveille S, Ferrucci L, Van Eijk JT, Guralnik JM. Exploring the effect of depression on physical disability: longitudinal evidence from the established populations for epidemiologic studies of the elderly. *Am J Public Health.* 1999; 89(9): 1346-1352.
- Begg S, Vos T, Barker B, Stevenson C, Stanley L, Lopez AD. The burden of disease and injury in Australia 2003. PHE 82. Canberra: AIHW; 2007.
- Tirupati S, Chua LE. Obesity and metabolic syndrome in a psychiatric rehabilitation service. *Aust N Z J Psychiatry.* 2007; 41(7): 606-10.
- Grundy SM, Brewer HB Jr, Cleeman JI, Smith SC Jr, Lenfant C. Definition of metabolic syndrome report of the National Heart, Lung, and Blood Institute/American Heart Association Conference on scientific issues related to definition. *Circulation.* 2004; 109(3): 433-8.
- Expert Panel on Detection Evaluation, and Treatment of High Blood Cholesterol in Adults. Executive summary of the Third Report of the National Cholesterol Education Program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III). *JAMA.* 2001; 285(19): 2486-97.
- Takeuchi T, Nakao M, Nomura K, Inoue M, Tsurugano S, Shinozaki Y, et al. Association of the metabolic syndrome with depression and anxiety in Japanese men: A 1-year cohort study. *Diabetes/metabolism research and reviews.* 2009; 25(8): 762-7.
- Pan A, Keum N, Okereke OI, Sun Q, Kivimaki M, Rubin RR, et al. Bidirectional association between depression and metabolic syndrome: a systematic review and meta-analysis of epidemiological studies. *Diabetes care.* 2012; 35(5): 1171-80.
- Hildrum B, Mykletun A, Midthjell K, Ismail K, Dahl AA. No association of depression and anxiety with the metabolic syndrome: the Norwegian HUNT study. *Acta Psychiatrica Scandinavica.* 2009; 120(1): 14-22.
- Whisman MA, Perez JE, Ramel W. Factor structure of the Beck Depression Inventory—Second Edition (BDI-II) in a student sample. *J Clin Psychol.* 2000; 56(4): 545-51.
- Ghassemzadeh H, Mojtabei R, Karamghadiri N, Ebrahimkhani N. Psychometric properties of a Persian-language version of the Beck Depression Inventory-Second edition: BDI-II-PERSIAN. *Depress Anxiety.* 2005; 21(4): 185-92.
- Aadahl M, Jørgensen T. Validation of a new self-report instrument for measuring physical activity. *Med Sci Sports Exerc.* 2003; 35(7): 1196-202.
- Kelishadi R, Rabiei K, Khosravi A, Famouri F, Sadeghi M, Rouhafza H, et al. Assessment of physical activity of adolescents in Isfahan. *J Shahrekord Univ Med Sci.* 2001. 3(2): 27-33.
- Tremblay AJ, Morrissette H, Gagné JM, Bergeron J, Gagné C, Couture P. Validation of the Friedewald formula for the determination of low-density lipoprotein cholesterol compared with  $\beta$ -quantification in a large population. *Clinical Chemists.* 2004; 37(9): 785-90.
- Hrebicek J, Janout V, Malinčíková J, Horáková D, Čížek L. Detection of insulin resistance by simple quantitative insulin sensitivity check index QUICKI for epidemiological assessment and prevention. *J Clin Endocrinol Metab.* 2002; 87(1): 144-7.
- Ra JS, Kim HS. Sex-based Association between Depression and Metabolic Syndrome in Korean



- Middle-aged and Older Adults. *Osong Public Health Res Perspect*. 2017; 8(2): 130-137.
23. Hiles SA, Révész D, Lamers F, Giltay E, Penninx BW. BIDIRECTIONAL PROSPECTIVE ASSOCIATIONS OF METABOLIC SYNDROME COMPONENTS WITH DEPRESSION, ANXIETY, AND ANTIDEPRESSANT USE. *Depress Anxiety*. 2016; 33(8): 754-64.
  24. Wiltink J, Michal M, Jünger C, Münzel T, Wild PS, Lackner KJ, et al. Associations between degree and sub-dimensions of depression and metabolic syndrome (MetS) in the community: results from the Gutenberg Health Study (GHS). *BMC Psychiatry*. 2018; 18: 114.
  25. Kahl KG, Greggersen W, Schweiger U, Cordes J, Balijepalli C, Löscher C, et al. Prevalence of the metabolic syndrome in unipolar major depression. *Eur Arch Psychiatry Clin Neurosci*. 2011; 262(4): 313-20.
  26. Dunbar JA, Reddy P, Davis-Lameloise N, Philpot B, Laatikainen T, Kilkkinen A, et al. Depression: an important comorbidity with metabolic syndrome in a general population. *Diabetes Care*. 2008; 31(12):2368-73.
  27. Olvera RL, Williamson DE, Fisher-Hoch SP, Vatcheva KP, McCormick JB. Depression, Obesity, and Metabolic Syndrome: Prevalence and Risks of Comorbidity in a Population-Based Study of Mexican Americans. *J Clin Psychiatry*. 2015; 76(10): e1300-5.
  28. Hussain T, Margoob MA, Shoib S, Shafat M, Chandel RK. Prevalence of Metabolic Syndrome among Psychiatric Inpatients: A Hospital Based Study from Kashmir. *J Clin Diagn Res*. 2017; 11(6): VC05-VC08.
  29. Takeuchi T, Nakao M, Nomura K, Yano E. Association of metabolic syndrome with depression and anxiety in Japanese men. *Diabetes Metab*. 2009; 35(1): 32-6.
  30. Pearson S, Schmidt M, Patton G, Dwyer T, Blizzard L, Otahal P, et al. Depression and insulin resistance: cross-sectional associations in young adults. *Diabetes Care*. 2010; 33(5): 1128-33.