

Effect of Intermittent Fasting during Ramadan on Visfatin, Adiponectin and Tumor Necrotizing Factor-Alpha in Healthy Muslim Individuals

Eisa Safavi¹, Ali Reza Rahbar^{2*}

1. Department of Laboratory Technology, Paramedical Faculty, Bushehr University of Medical Sciences, Bushehr, Iran
2. Department of Nutrition, Persian Gulf Tropical Medicine Research Center, Bushehr University of Medical Sciences, Bushehr, Iran

ARTICLE INFO

Article type:
Original article

Article History:
Received: 27 Mar 2017
Accepted: 21 May 2017
Published: 20 Jun 2017

Keywords:
Adiponectin
Ramadan fasting
Tumor necrotizing factor-
alpha
Visfatin

ABSTRACT

Introduction: The aim of this study was to investigate the effect of fasting in Ramadan on visfatin, tumor necrosis factor alpha (TNF- α) and adiponectin level in human.

Methods: Thirty four men aged 24 to 55 years old were selected from those people who were willing to fast in Ramadan. The blood sample was obtained from each participant in fasting state at the beginning and the end of study to determine serum visfatin, adiponectin and TNF- α . Paired T test was used to identify differences between beginning and the end of the study in serum visfatin, adiponectin and TNF- α .

Results: Results showed a significant decrease in Visfatin after study. There were no significant changes in adiponectin and TNF- α during Ramadan fasting.

Conclusion: It is concluded that fasting in Ramadan is beneficial to health and can ameliorate some inflammatory markers in fasting individuals.

► Please cite this paper as:

Safavi E, Rahbar AR. Effect of Intermittent Fasting during Ramadan on Visfatin, Adiponectin and Tumor Necrotizing Factor- Alpha in Healthy Muslim Individuals. J Fasting Health. 2017; 5(2): 50-55. DOI: [10.22038/jfh.2017.22719.1084](https://doi.org/10.22038/jfh.2017.22719.1084)

Introduction

Inflammation is primitive in athero-sclerosis (1) and plays an essential role in the development of CVD (2). Recently, growing interest has appeared in the role of inflammatory and anti-inflammatory adipocytokines in the progression of CVD (3, 4). Visfatin, as an inflammatory adipocytokine, is involved in many inflammatory diseases (5). Reportedly, Visfatin has been engaged in monocyte adhesion to endothelial cells, (6) and plaque rupture (7) in unstable lesions in atherosclerosis. On the contrary, adiponectin is an anti-inflammatory adipocytokine, (8) which prevents vascular stenosis, (9) insulin resistance, atherosclerosis, (10) renal fibrosis, diabetes nephropathy and albuminuria in animal models (10). In addition, TNF- α induces

programmed cell death or apoptosis in Rheumatoid arthritis (RA), (11) Systemic Lupus Erythematosus (SLE) (12) and Ankylosing spondylitis (AS) (13).

Evidences show that fasting is effective in the treatment of rheumatoid diseases, and chronic pain syndromes (14). Fasting is used by Naturopathy to manage treatment of diseases (15). Ayurveda also use fasting to enhance digestive capacity, enthusiasm, and decrease signs and symptoms of diseases (16). However, there are controversies about the period and the rate of calorie restriction in different studies (17). Some studies recommend severe restriction, up to 75-90% of energy needs (18). But prolonged and severed starvation has hazardous effects such

* Corresponding author: Ali Reza Rahbar, Department of Nutrition, Persian Gulf Tropical Medicine Research Center, Bushehr University of Medical Sciences, Bushehr, Iran. Tel: 00989177724604; Fax: + 0098 7712528; Email: rahbar_alireza@yahoo.com

© 2017 mums.ac.ir All rights reserved.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

as heart failure, (19) impaired liver function, (20) cholesterogenesis, (21) autophagy (21) and cachexia (22). It has been hypothesized that intermittent fasting positively affects the inflammatory markers (23). We hypothesize that intermittent fasting might be able to ameliorate some health-threatening factors such as elevated TNF- α or Visfatin without the undesirable side effects of prolonged starvation. However, to the best of our knowledge, so far, there is very scarce published study on the effect of intermittent fasting on visfatin and TNF- α during Ramadan. Therefore this study was conducted to investigate the effect of fasting in Ramadan on visfatin, TNF- α and adiponectin level in human.

Material and methods

This study was conducted between 1st and 30th of July in 2015 in Bushehr University of Medical Sciences. Thirty four men aged 16 to 64 years old were selected by simple random sampling from those people who were willing to participate in the study. The participants were invited to present in Bushehr University of Medical Sciences between 8-9 hours on the following morning in fasting state through a letter delivered to their homes by the post. The aims of the study were elucidated in detail in the letter. The inclusion criteria were: willing to fast during Ramadan, male gender and aged 15 years old and more (eligibility criteria for fasting from the religious point of view for male gender). They were excluded if they were using thyroid drug, cigarette, estrogen, medications for hypertension and any clinical signs of hypothyroidism, dyslipidemia, or diabetes mellitus. An expert physician and dietitian interviewed and examined the participants.

At the beginning of study a blood sample was obtained from each participant after 12 hours fasting into tube containing EDTA. Blood sample were centrifuged at 3000 \times g for 15 min and the plasma samples were stored in 1.5 ml microtubes at -20 $^{\circ}$ C at the laboratory of Persian Gulf Tropical Medicine Research. At the end of the study once again plasma samples were obtained from each participant and plasma

visfatin, TNF- α and adiponectin were measured. A written, signed informed consent was obtained from each participant at the beginning of the study.

Anthropometric measurements

Height and weight of the participants were measured with a stadiometer (Seca stadiometer, model 286 dp, Hamburg, Germany) and a scale (Seca scale, model 755, Hamburg, Germany).

The participants were demanded to remove heavy outer garments and shoes before height and weight measurement. Body mass index was calculated as weight in kilograms divided by the square of the height in meters (kg/m²). Waist circumference was measured at the midpoint between the lower edge of the rib cage and the iliac crests with an un-stretchable measuring tape. Waist-to-hip ratio was determined as waist circumference divided by hip circumference. The smallest circumference measured at the navel, and the largest circumference measured at the hips and buttocks was applied for this measurement (24).

Physical activity

Physical activity of participants was assessed by a questionnaire based on the International Physical Activity Questionair, 2005, Epic-Norfolk.

Dietary intake analysis

Trained dietitians recorded a 24-hour dietary recall for 3 nonconsecutive days from each participant before and during Ramadan. Nutritionist III software (version 7.0; NSquared Computing, Salem, OR, USA) were used to analyze the content of energy and other nutrients of each food and beverage. This software was adapted for Iranian foods.

The protocol of the study was endorsed by the research deputy of Bushehr University of Medical Sciences (DP/8703277/176, 14/4/2013). The ethics committee of Bushehr University of Medical Sciences (May 2015) approved the ethical aspects of the study, and the research reported here was carried out in accordance with the principles of the Declaration of Helsinki as revised in 2000.

Measurement of serum parameters

Plasma Visfatin was measured by a

commercially available ELISA kit (Bioassay Technology Laboratory, Shanghai crystal day biotech co, LTD, Shanghai, China, Catalog No. E0025Hu). The assay range was 0.5 ng/mL-100ng/mL, and the intra- and inter-assay coefficients of variance were less than 10% and less than 12%, respectively.

To detect Adiponectin in plasma samples, a commercially available ELISA kit (Bioassay Technology Laboratory, Shanghai crystal day biotech co, LTD, Shanghai, China, Catalog No. E1550Hu) was used according to the manufacturer's instructions. The assay range was 0.2 mg/L-60mg/L, and the intra- and inter-assay coefficients of variance were less than 10% and less than 12%, respectively.

Plasma TNF- α was measured by a commercially available ELISA kit (Bioassay Technology Laboratory, Shanghai crystal day biotech co, LTD, Shanghai, China, Catalog No. E0082Hu). The assay range was 3 ng/L-1.52ng/L, and the intra- and inter-assay coefficients of variance were less than 10% and less than 12%, respectively.

Statistical analysis

The distribution of variables was studied with probability plots and the Shapiro-Wilks test. Paired T test was used to identify differences between beginning and the end of the study in serum visfatin, adiponectin and TNF- α .

The relationship between variables was determined by calculating Pearson's correlation coefficient. A value of $p < 0.05$ was accepted as significant.

All statistical analyses were done with an

IBM computer and the SPSS v. 15 statistical software packages (SPSS Inc., Chicago, IL, USA) were used to analyze the data statistically.

Results

The characteristics of the participants are shown in Table 1. Bivariate correlation analysis showed no significant correlation between change in serum visfatin, adiponectin and TNF- α level and change in WHR and BMI or calorie intake but there was a significant association between serum visfatin, adiponectin and TNF- α level and age in population study (data not shown).

The paired sample T- test showed 11.79 % significant decrease in Visfatin during Ramadan fasting. TNF- α and adiponectin did not change significantly after Ramadan fasting compared to before Ramadan. (Table 2).

The paired sample T- test showed a 0.5 % significant decrease in weight and waist circumferences during Ramadan fasting. (Table 2).

The energy, carbohydrate and fat intake

Table 1. Characteristics of the individuals (n=34). Means \pm SD

Characteristic NO	individuals	
	(n=34) mean	SD
Age(years)	35	11
BMI (kg/m ²)	24.79	3.08
WHR	0.908	0.109
weight (Kg)	74.62	10.63
Waist circumference (Cm)	91.97	11.71
Visfatin (ng/mL)	42.58	21.44
Adiponectin (mg/L)	22.30	10.43
TNF- α (ng/L)	289.92	181.22

BMI: body mass index, WHR: Waist Hip Ratio, igf-1: Insulin like growth hormone-1, TNF- α : Tumor necrosis factor-alpha

Table 2. Change in variables before and after Ramadan unadjusted and age adjusted

variables	Before Ramadan	After Ramadan	Unadjusted P value	Adjusted P value
Visfatin(ng/mL)	42.58 \pm 21.44	40.17 \pm 20.44	0.045	0.047
Adiponectin(mg/L)	22.30 \pm 10.43	21.48 \pm 10.50	0.14	0.23
TNF- α (ng/L)	289.92 \pm 181.22	277.89 \pm 180.58	0.39	0.55
Weight(kg)	74.62 \pm 10.63	73.93 \pm 10.25	0.094	0.13
BMI(kg/m ²)	24.79 \pm 3.08	24.57 \pm 3.05	0.11	0.42
WC(Cm)	91.97 \pm 11.71	91.45 \pm 11.59	0.029	0.032
WHR	0.908 \pm 0.109	0.910 \pm 0.115	0.72	0.94
Energy(Calorie)	2392.06 \pm 664.48	1671.35 \pm 568.13	0.000	0.001
Fat (gm/day)	62.54 \pm 22.43	33.34 \pm 23.03	0.000	0.001
SFA (gm/day)	34.65 \pm 16.35	16.36 \pm 15.85	0.000	0.001
MUFA(gm/day)	14.20 \pm 5.34	9.37 \pm 6.483	0.003	0.013
PUFA(gm/day)	11.93 \pm 11.31	5.57 \pm 10.21	0.034	0.034
Carbohydrate (gm/day)	377.51 \pm 141.58	302.90 \pm 106.95	0.01	0.03
Protein(gm/day)	87.62 \pm 39.87	51.08 \pm 19.16	0.000	0.002

BMI: body mass index, WHR: Waist Hip Ratio, WC: Waist circumference, TNF- α : Tumor necrosis factor-alpha, SFA: Saturated fatty acids, PUFA: Poly unsaturated fatty acids, MUFA: Mono unsaturated fatty acid

decreased significantly 30.12 %, 19.76% and 46.69 % respectively (Table 2).

Discussion

The data of present study showed a significant decrease in visfatin level in individuals who fasted at least 16 hours per day for one month. Studies with severe fasting failed to change visfatin level in human model (25, 26). Owczarek et al, (2016) supported the existence of an association between nutritional status, inflammation and circulating visfatin level (27). This relationship is in connection with the change of leptin concentration after intermittent fasting (28). Low concentration of leptin is prevalent in energy deprivation in humans (29, 30). The adverse relationship between Leptin and visfatin level is addressed by Tan and et al (31). Chowdhury and et al, (2016) reported a significant lower Systemic concentration of leptin in the afternoon following morning fasting (32). A significant decrease in blood leptin also is reported in fasted individuals during Ramadan (33).

In this study no significant changes were observed in TNF- α sequel fasting. Although in animal model significant decrease were seen after intermittent and prolonged fasting, (34, 35) intermittent fasting in human in correspondence to the present study did not affect the TNF- α (36). It should be mentioned that Ramadan fasting is accompanied by alterations in meal times, meal composition, sleep/wake patterns, circadian rhythms and physical activity (37). Ramadan fasting as a stressor stimulates stress responses in individuals (38). Psychological and physiological stress induces inflammatory markers (39). And this might influence the beneficial effect of intermittent fasting on TNF- α .

The concentration of anti-inflammatory cytokine (adiponectin) was not changed after intermittent fasting during this study. Yannakoulia and colleagues, (2003) in agreement with our findings reported no significant relationship between calorie intake and serum adiponectin level (40). In fact, the concentration of adiponectin compared to other cytokines in the circulation may be too high to be influenced by alteration in intermittent fasting. It has been reported that plasma levels of adiponectin range from 5 to 30 mg/ L in

normal weight individuals representing 0.01% of all plasma protein (41) and up to 1000 fold greater than visfatin (Table 1). Furthermore, in animal studies which have demonstrated a significant increase in adiponectin level following intermittent fasting, a calorie restriction was applied concomitant with fasting (42). However, in current study the participant were free to eat after fasting.

Conclusion

The findings of this study revealed that intermittent fasting independent of changes in anthropometric measures decreases visfatin level in human.

Conflicts of interest

We performed this research in the interest of knowledge and declare that we have no conflict of interest.

Acknowledgment

This study was supported in part by a grant from Bushehr University of Medical Sciences, Bushehr, Iran (DP/20/ 18/3/2316, 1/1/2015). We thank K. Shashok (Author AID in the Eastern Mediterranean) for improving the use of English in the manuscript.

References

1. Ross R. Atherosclerosis--an inflammatory disease. *N Engl J Med.* 1999; 340(2):115-26.
2. Ribeiro F, Alves AJ, Duarte JA, Oliveira J. Is exercise training an effective therapy targeting endothelial dysfunction and vascular wall inflammation? *Int J Cardiol.* 2010; 141(3):214-21.
3. Bulcão C, Ferreira SR, Giuffrida FM, Ribeiro-Filho FF. The new adipose tissue and adipocytokines. *Curr Diabetes Rev.* 2006; 2(1):19-28.
4. Yang RZ, Lee MJ, Hu H, Pray J, Wu HB, Hansen BC, et al. Identification of omentin as a novel depot-specific adipokine in human adipose tissue: possible role in modulating insulin action. *Am J Physiol Endocrinol Metab.* 2006; 290(6):E1253-61.
5. Chandran M, Phillips SA, Ciaraldi T, Henry RR. Adiponectin: more than just another fat cell hormone? *Diabetes Care.* 2003; 26(8):2442-50.
6. Kim SR, Bae YH, Bae SK, Choi KS, Yoon KH, Koo TH, et al. Visfatin enhances ICAM-1 and VCAM-1 expression through ROS-dependent NF- κ B activation in endothelial cells. *Biochim Biophys Acta.* 2008; 1783(5):886-95.
7. Dahl TB, Yndestad A, Skjelland M, Øie E, Dahl A, Michelsen A, et al. Increased expression of visfatin in macrophages of human unstable carotid and

- coronary atherosclerosis possible role in inflammation and plaque destabilization. *Circulation*. 2007; 115(8):972-80.
8. Goldstein BJ, Scalia R. Adiponectin: a novel adipokine linking adipocytes and vascular function. *J Clin Endocrinol Metab*. 2004; 89(6):2563-8.
 9. Matsuda M, Shimomura I, Sata M, Arita Y, Nishida M, Maeda N, et al. Role of adiponectin in preventing vascular stenosis. The missing link of adipo-vascular axis. *J Biol Chem*. 2002; 277(40):37487-91.
 10. Yamauchi T, Kamon J, Waki H, Imai Y, Shimozawa N, Hioki K, et al. Globular adiponectin protected ob/ob mice from diabetes and ApoE-deficient mice from atherosclerosis. *J Biol Chem*. 2003; 278(4):2461-8.
 11. Barton A, John S, Ollier WE, Silman A, Worthington J. Association between rheumatoid arthritis and polymorphism of tumor necrosis factor receptor II, but not tumor necrosis factor receptor I, in Caucasians. *Arthritis Rheum*. 2001; 44(1):61-5.
 12. Studnicka-Benke A, Steiner G, Petera P, Smolen JS. Tumor necrosis factor alpha and its soluble receptors parallel clinical disease and autoimmune activity in systemic lupus erythematosus. *Rheumatology*. 1996; 35(11):1067-74.
 13. Bal A, Unlu E, Bahar G, Aydog E, Eksioglu E, Yorgancioglu R. Comparison of serum IL-1 β , sIL-2R, IL-6, and TNF- α levels with disease activity parameters in ankylosing spondylitis. *Clin Rheumatol*. 2007; 26(2):211-5.
 14. Michalsen A, Li C. Fasting therapy for treating and preventing disease-current state of evidence. *Forsch Komplementmed*. 2013; 20(6):444-53.
 15. Nair PM, Nanda A. Naturopathic medicine in India. *Focus Alternat Complement Ther*. 2014; 19(3):140-7.
 16. Shripathi Adiga H, Adiga RS. Concept and canons of fasting in Ayurveda. *J Fast Health*. 2013; 1(1):37-40.
 17. Fontana L, Weiss EP, Villareal DT, Klein S, Holloszy JO. Long-term effects of calorie or protein restriction on serum IGF-1 and IGFBP-3 concentration in humans. *Aging Cell*. 2008; 7(5):681-7.
 18. Harvie MN, Pegington M, Mattson MP, Frystyk J, Dillon B, Evans G, et al. The effects of intermittent or continuous energy restriction on weight loss and metabolic disease risk markers: a randomized trial in young overweight women. *Int J Obes*. 2011; 35(5):714-27.
 19. Grossniklaus DA, O'Brien MC, Clark PC, Dunbar SB. Nutrient intake in heart failure patients. *J Cardiovasc Nurs*. 2008; 23(4):357.
 20. Elliott B, Mina M, Ferrier C. Complete and voluntary starvation of 50 days. *Clin Med Insights Case Rep*. 2016; 9:67-70.
 21. Godavarthy P, Kumari YS, Bikshapathy E. Starvation induced cholesterologenesis in hepatic and extra hepatic tissues of climbing Perch, *Anabas testudineus* (Bloch). *Saudi J Biol Sci*. 2012; 19(4):489-94.
 22. Evans WJ, Morley JE, Argilés J, Bales C, Baracos V, Guttridge D, et al. Cachexia: a new definition. *Clin Nutr*. 2008; 27(6):793-9.
 23. Faris MA, Kacimi S, Al-Kurd RA, Fararjeh MA, Bustanji YK, Mohammad MK, et al. Intermittent fasting during Ramadan attenuates proinflammatory cytokines and immune cells in healthy subjects. *Nutr Res*. 2012; 32(12):947-55.
 24. Consultation WE. Waist circumference and waist-hip ratio. Report of a WHO Expert Consultation. Geneva: World Health Organization; 2008. P. 8-11.
 25. Mitterberger MC, Mattesich M, Klaver E, Lechner S, Engelhardt T, Larcher L, et al. Adipokine profile and insulin sensitivity in formerly obese women subjected to bariatric surgery or diet-induced long-term caloric restriction. *J Gerontol A Biol Sci Med Sci*. 2010; 65(9):915-23.
 26. Kang ES, Magkos F, Sienkiewicz E, Mantzoros CS. Circulating vaspin and visfatin are not affected by acute or chronic energy deficiency or leptin administration in humans. *Eur J Endocrinol*. 2011; 164(6):911-7.
 27. Owczarek AJ, Olszanecka-Glinianowicz M, Kocelak P, Bozentowicz-Wikarek M, Brzozowska A, Mossakowska M, et al. The relationship between circulating visfatin/nicotinamide phosphoribosyltransferase, obesity, inflammation and lipids profile in elderly population, determined by structural equation modeling. *Scand J Clin Lab Invest*. 2016; 76(8):632-40.
 28. Garcia-Fuentes E, Garcia-Almeida JM, Garcia-Arnes J, Garcia-Serrano S, Rivas-Marin J, Gallego-Perales JL, et al. Plasma visfatin concentrations in severely obese subjects are increased after intestinal bypass. *Obesity (Silver Spring)*. 2007; 15(10):2391-5.
 29. Dardeno TA, Chou SH, Moon HS, Chamberland JP, Fiorenza CG, Mantzoros CS. Leptin in human physiology and therapeutics. *Front Neuroendocrinol*. 2010; 31(3):377-93.
 30. Rosenbaum M, Nicolson M, Hirsch J, Murphy E, Chu F, Leibel RL. Effects of weight change on plasma leptin concentrations and energy expenditure. *J Clin Endocrinol Metab*. 1997; 82(11):3647-54.
 31. Tan BK, Chen J, Brown J, Adya R, Ramanjaneya M, Menon V, et al. In vivo and ex vivo regulation of visfatin production by leptin in human and murine adipose tissue: role of mitogen-activated protein kinase and phosphatidylinositol 3-kinase signaling pathways. *Endocrinology*. 2009; 150(8):3530-9.
 32. Chowdhury EA, Richardson JD, Tsintzas K, Thompson D, Betts JA. Effect of extended morning fasting upon ad libitum lunch intake and associated metabolic and hormonal responses in obese adults. *Int J Obes*. 2016; 40(2):305-11.

33. Khoshdel A, Kheiri S, Nasiri J, Tehran HA, Heidarian E. The effect of Ramadan fasting on serum leptin, neuropeptide Y and insulin in pregnant women. *Med J Islam Repub Iran*. 2014; 28:92.
34. Razeghi Jahromi S, Ghaemi A, Alizadeh A, Sabetghadam F, Moradi Tabriz H, Togha M. Effects of intermittent fasting on experimental autoimmune encephalomyelitis in C57BL/6 mice. *Iran J Allergy Asthma Immunol*. 2016; 15(3):212-9.
35. Vasconcelos AR, Yshii LM, Viel TA, Buck HS, Mattson MP, Scavone C, et al. Intermittent fasting attenuates lipopolysaccharide-induced neuroinflammation and memory impairment. *J Neuroinflammation*. 2014; 11:85.
36. Feizollahzadeh S, Rasuli J, Kheirouri S, Alizadeh M. Augmented plasma adiponectin after prolonged fasting during Ramadan in men. *Health Promot Perspect*. 2014; 4(1):77-81.
37. Bahammam A. Does Ramadan fasting affect sleep? *Int J Clin Pract*. 2006; 60(12):1631-7.
38. Lahdimawan A, Handono K, Indra MR, Prawiro SR. Effect of Ramadan fasting on endorphin and endocannabinoid level in serum, PBMC and macrophage. *Int J Pharm Sci Invent*. 2013; 2(3):46-54.
39. Lin J, Shen Y, Gao Y, Li L. beta-Endorphin enhances IL-2 and IFN-gamma gene expression in human blood mononuclear cells. *Zhongguo Yi Xue Ke Xue Yuan Xue Bao*. 1997; 19(5):353-6.
40. Yannakoulia M, Yiannakouris N, Blüher S, Matalas AL, Klimis-Zacas D, Mantzoros CS. Body fat mass and macronutrient intake in relation to circulating soluble leptin receptor, free leptin index, adiponectin, and resistin concentrations in healthy humans. *J Clin Endocrinol Metab*. 2003; 88(4):1730-6.
41. Lihn AS, Bruun JM, He G, Pedersen SB, Jensen PF, Richelsen B. Lower expression of adiponectin mRNA in visceral adipose tissue in lean and obese subjects. *Mol Cell Endocrinol*. 2004; 219(1):9-15.
42. Wan R, Ahmet I, Brown M, Cheng A, Kamimura N, Talan M, et al. Cardioprotective effect of intermittent fasting is associated with an elevation of adiponectin levels in rats. *J Nutr Biochem*. 2010; 21(5):413-7.