



Nutritional Assessment and Dietary Requirements in Kidney Transplant Patients: A Literature Review as a Clinical Guideline

Sudiyeh Hejri Zarifi¹, Naseh Pahlavani^{1, 2}, Farveh Yahyapour¹, Mohsen Nematy¹, Abdolreza Norouzy^{1*}

1- Department of Nutrition, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran.

2- Student Research Committee, Mashhad University of Medical Sciences, Mashhad, Iran.

ARTICLE INFO	ABSTRACT
<i>Article type:</i> Review Article	Kidney transplantation is a preferred treatment and the 'last resort' in chronic kidney disease after end-stage renal disease. Considering the key role of nutrition in post-transplant recovery, the present study aimed to review the nutritional assessment and dietary requirements of kidney transplant patients. Relevant articles were retrieved via searching in databases such as PubMed, Medline, Google Scholar, Scopus, and Web of Science using various keywords and phrases, including kidney transplantation, end-stage disease, nutrition, dietary intake, nutritional assessment, and nutritional evaluation. According to the literature, medical nutrition therapy is essentially involved in the post-transplantation recovery of patients with kidney failure. Furthermore, it seems that nutritional and medical evaluation based on laboratory methods, vital signs, and anthropometric measurements are critical to improving the quality of nutritional interventions after kidney transplantation and may help decrease the risk of kidney transplant rejection. Therefore, the periodic evaluation and follow-up of these could yield beneficial outcomes.
<i>Article History:</i> Received: 26 Aug 2020 Accepted: 20 Sep 2020 Published: 17 May 2021	
<i>Keywords:</i> Kidney Transplantation Nutritional Assessment Dietary Requirements Food Intake	
► <i>Please cite this paper as:</i> Hejri Zarifi S, Pahlavani N, Yahyapour F, Nematy M, Norouzy A. Nutritional Assessment and Dietary Requirements in Kidney Transplant Patients: A Literature Review as a Clinical Guideline. <i>J Nutr Fast Health.</i> 2021; 9(2): 113-119. DOI: 10.22038/JNFH.2020.51553.1292	

Introduction

Chronic kidney disease (CKD) is a general term referring to various disorders that affect the structure and function of the kidneys. One of the definitions of CKD is the reduction of the glomerular filtration rate to less than 60 ml/min per 1.73 m² of the body surface (1). The incidence of CKD varies in different regions across the world. In general, the incidence of the disease in most countries has been reported to be higher than 200 cases per million annually (2).

Kidney transplantation is the preferred treatment and the 'last resort' for CKD after end-stage renal disease (ESRD) (3). In kidney transplantation, nutritional requirements change entirely due to comorbidities, immunosuppressive drugs, and recovery from major surgery, thereby leading to malnutrition and mineral deficiencies. Statistics suggest that more than 70% of CKD patients experience malnutrition (4, 5). Several factors may contribute to the deterioration of nutritional status in these patients, including obesity, renal

replacement therapy, anorexia, and dietary restrictions. Therefore, the periodic evaluation of nutritional status in these patients by a clinical nutritionist is essential (6, 7).

A proper dietary pattern may reduce the risk of transplant rejection and length of hospital stay and prevent threatening diseases, obesity, and multiple adverse conditions associated with transplants (8). According to a study in this regard, controlling the nutritional status and providing a proper and healthy diet to kidney transplant recipients could reduce body fat and body weight, while also contributing to an ideal weight and improved cholesterol, triglyceride, and blood sugar levels (9). The nutritional assessment of kidney transplant recipients during the treatment process could prompt timely and appropriate interventions. Among the most important nutritional assessment tools are subjective global assessment (SGA) forms, immunological tests, and biochemical and anthropometric indices, which could be largely

* Corresponding author: Abdolreza Norouzy, Associate Professor, Department of Nutrition, Faculty of Medicine, Mashhad University of Medical Sciences, Azadi Sq., Mashhad, Iran; Tel: +985138002382, Email: norouzya@mums.ac.ir.

© 2021 mums.ac.ir All rights reserved.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

beneficial for the nutritional status monitoring of these patients (10).

This comprehensive review aimed to evaluate the nutritional assessment methods used for kidney transplant recipients, as well as the nutritional requirements of these patients based on the most recent guidelines and studies.

Materials and Methods

Search Strategy

Relevant articles were identified via searching in databases such as PubMed, Medline, Scopus, and Web of Science databases using various keywords, including "kidney transplantation", OR "kidney transplant", OR "end-stage renal disease", ", OR "renal transplant" OR "kidney failure" OR "renal failure" in combination with "nutrition" "dietary requirements", "nutrition intervention", "nutritional assessment", "nutrition evaluation", "dietary assessment", "food intake", and "nutrition intakes". The identified studies and review articles were assessed, and eligible studies were selected for the review, including reports on the nutritional and dietary requirements of kidney transplantation. Notably, our literature review was not performed within a systematic framework, and duplicates and unrelated articles were excluded.

Nutritional Assessments Tools

Candidates for kidney transplantation often have an abnormal nutritional status due to organ dysfunction and the associated symptoms (11). The importance of the nutritional assessment of kidney transplant recipients is due to the need for an initial and general assessment, followed by detailed examination (12). Several methods are applied to assess the nutritional status of kidney transplant recipients, some of which have been discussed in the following sections.

Medical History and Social Status

Medical history and social status encompass primary and secondary diagnosis, disease history, previous surgeries, previous weight, history of weight changes, and health assessment of various organs (gastrointestinal tract, liver and bile ducts, circulation, cardiovascular system, nerves, lungs, and immune system). Social status refers to the data on socioeconomic status, religion, living facilities, and food security (13). The most common clinical symptoms in kidney transplant recipients include edema, fatigue, nausea and vomiting, and anorexia, which should be evaluated meticulously (14).

Anthropometric Indices

Anthropometry is the concept of measuring the body size and the related scales. Anthropometric measurements indicate health and the growth status and are commonly used to assess nutritional status, such as obesity due to overeating and weight loss due to protein-energy malnutrition (13, 15). These data are beneficial in determining the impact of nutritional interventions on various diseases, traumas, surgeries or malnutrition. Anthropometrics are also a method used for the estimation of the body mass in clinical trials (16). Height and weight measurements are among key anthropometric indicators, and body mass index is employed as an indirect method to determine obesity or body fat percentage (16). If height and weight cannot be measured (e.g., in unconscious or bedridden patients) to estimate the muscle mass (body protein stores) and fat stores, waist circumference (WC) and mid-arm muscle circumference are the alternatives. WC is an estimate of the subcutaneous and intra-abdominal distribution of the adipose tissue, as well as a simple anthropometric index associated with the visceral adipose tissue. If WC is higher than 102 centimeters in men and higher than 88 centimeters in women, it indicates the increased risk of disease and mortality (17).

Although previous findings have introduced an association between the primary grade of obesity with increased survival in ESRD patients, the hypothesis has been ruled out, and a normal body mass in these patients has been associated with better outcomes (18). In addition, a normal body mass has been associated with improved quality of life and reduced complications, such as wound infections, delayed graft function, and rejection in kidney transplant recipients (19, 20). The measurement of triceps skinfold thickness is also used to measure the subcutaneous fat reserves of the body (21).

Bioelectrical Impedance Analysis (BIA)

Bioelectrical impedance analysis (BIA) is a practical method to measure the body composition (22). BIA is commonly used for estimating the body composition, especially the body water, body fat, and muscle mass (22). The BIA device measures parameters such as resistance and reactance, with the former showing the hydration status of the patient in association with body water and the latter

indicating the available energy to the body tissues (capacitator) (12, 23).

Food Intake Assessment

Nutritional intake is an inherent element of nutritional assessment as these evaluations provide data on the quantity and quality of the diet, appetite changes, food sensitivity, food intolerance, and causes of inadequate food intake during illness or in the recovery period (13). The food frequency questionnaire (FFQ) is a primary tool used for dietary assessment, collecting data on the proper amount and consumption of foods (per week, month or year). Furthermore, FFQ is a retrospective evaluation of food consumption frequency (24). The 24-hour recall and dietary record are the other methods of assessing food intake in these patients, which accurately measure their dietary intake (25, 26).

SGA

The SGA consists of two sections, which are medical history and physical examination. The first section is based on the medical history of the patient and has five components, including weight changes, diet, gastrointestinal symptoms, functional capacity, and underlying hypercatabolic diseases (e.g., physical injuries, burns, inflammatory diseases, infections, and malignancies) (27-29). The second section of the SGA is physical examination, which has the three components of subcutaneous fat analysis, muscle analysis, and the presence of edema/ascites. If these components are normal, they are defined as component A. If they are mildly to moderately affected, they are rated B, and if they are severely affected, they are graded C (27). In the final scoring of the SGA, if most of the components are scored A, the patient has a normal nutritional status, and if most of the components are scored B, mild-to-moderate malnutrition is confirmed. If

most of the components are scored C, severe malnutrition is recognized (27, 28).

Evaluation of Biochemical Data

Biochemical tests are highly sensitive and could be altered by food/medications or certain diseases. Blood and urine are the main sources of biochemical tests (30). Visceral protein measurement includes the measurement of total serum proteins (albumin and transferrin), the changes in which occur due to dietary protein deficiency, altered metabolism and body fluid status, inflammation, and use of certain medications (30, 31). Among the other biochemical measurements are creatinine, serum levels of sodium, potassium, calcium, magnesium, and phosphorous, fasting blood sugar, liver enzymes, blood urea nitrogen, and lipid profile, which play a key role in the monitoring and management of the clinical and nutritional status of patients with CKD (30, 32).

Nutritional Requirements of Kidney Transplantation

In kidney transplantation, the patient has a wider choice of diets and food intake and often require less medical, nursing, and nutritional care compared to hemodialysis patients (Table 1) (33). The nutritional care of kidney transplant recipients normally depends on the metabolic effects of treatment with immunosuppressive drugs, such as corticosteroids, cyclosporine, and azathioprine (34). However, the side-effects of anti-rejection medications (e.g., diarrhea, constipation, nausea and vomiting, high glucose, high potassium [cyclosporine]) may occur after transplantation and affect food intake, thereby requiring medical and nutritional interventions (35).

Table 1. Comparison between hemodialysis and kidney transplantation

Variable	Hemodialysis patients	Transplant patients
Primary Treatment responsibility	Health care personnel	Patient
Diet	Low K, low PO ₄ , low Na, moderate protein, fluid restriction	High and moderate protein, no K or PO ₄ restrictions, no fluid restriction
Location	Clinical dialysis unit	No limitations
Risks	Bleeding, sepsis, infection	Immuno-compromised, diabetes, cancer
Contraindications	Poor cardiac status, poor blood vessels for access	High body mass index, noncompliance with medications

Corticosteroid intakes has been reported to increase protein catabolism, hyperlipidemia, sodium retention, weight gain, and glucose intolerance, while also inhibiting the normal metabolism of calcium, phosphorus, and vitamin

D (36-38). Cyclosporine therapy is associated with hyperkalemia, hyperlipidemia, and hypertension; therefore, it is essential to control the nutritional status of the recipients (39). The control of nutritional status (anthropometric

assessments, food intake, and food allergies) and routine biochemical tests are also important in this regard, and these patients should be periodically monitored by nutritionists (40, 41). The dietary pattern following kidney transplantation is often divided into two stages, which are the first month (4-6 weeks) after transplantation and more than one month after transplantation. During the first month, a high-protein diet (1.3-1.5 g/kg/day) with the energy level of 30-35 kilocalories is recommended to prevent negative nitrogen balance, while more protein (1.6-2 g/kg/day) is usually required in the patients with infections, surgery or stress (4, 39).

The protein recommendations in the chronic post-transplant population may vary depending on the diabetes status. A patient without diabetes mellitus (DM) may have an estimated protein requirement of 0.6-0.8 g/kg/day, while a patient

diagnosed with DM might have slightly higher protein requirements (0.9-1 g/kg/day) (Table 2) (42). Fat intake recommendations in the acute and chronic phase of transplantation are 25-30% and 30-35% of the total calories, respectively. It is important to emphasize on the consumption of monounsaturated and polyunsaturated fatty acids, minimal intake of trans fats, and low intake of saturated fats (43). According to the National Kidney Foundation, carbohydrates should provide 50-65% of the non-protein calories in the acute post-transplant phase, while in the chronic post-transplant phase, they should be reduced to 45-50% of the total energy (43). In addition, sodium restriction to 80-100 mEq/day minimizes fluid retention and contributes to blood pressure control. After one month, protein intake could be reduced to 1 g/kg/day, and sodium intake is adjusted individually (39).

Table 2. Nutritional requirements in kidney transplantation

	Acute phase after kidney transplantation (Less than 1-month)	Chronic phase after kidney transplantation (More than 1-month)
Energy (Kcal)	Calculate with the formula provided at the bottom of the table * or (30-35 kcal/kg IBW**)	Calculation with pre-dialysis formula (32-35 kcal/kg IBW**)
Protein (g/Kg/day)	Transplant from living person 1.3-1.5 Transplant from brain death person; 1.1-1.3	Without DM; 0.6-0.8 g/Kg/day With DM; 0.9-1 g/Kg/day
High biological value (HBV) protein	50% of total protein	50% of total protein
Fat	25-30% from total calorie	30-35% from total calorie
Carbohydrate	Remaining percentage of calories (50-65% of nonprotein calories)	Remaining percentage of calories (45-50% of total energy)
Potassium	Without limitation	Without limitation
Phosphorus	Without limitation	Without limitation
Sodium	2000-3000 mg/day	3000-4000 mg/day
Fluids	500-600 ml+ Excreted urine volume+ Excretion of urine from other routes	Without limitation
Medication side effects	Adjust diet to address the potential for hyperglycemia and hyperkalemia; observe food safety practices to minimize the risk for food-borne infections	
Bone health	Monitor laboratory values. Consider need for supplementation of calcium, magnesium, phosphorus, and vitamin D	

* Total daily energy requirements in the acute phase of kidney transplantation=[Basal metabolic rate × Stress Coefficient × Physical activity coefficient].

Stress Coefficient in kidney surgery=1.2

Stress Coefficient in Inpatients= 1.2

Stress Coefficient in Outpatients= 1.3

Abbreviations: IBW; Ideal Body Weight, DM; Diabetes Mellitus

Hyperkalemia is associated with cyclosporine intake and often requires potassium restriction although it may be temporary (34). After kidney transplantation, most patients present with mild hypophosphatemia and hypercalcemia due to the bone resorption induced by hyperparathyroidism and the effects of

corticosteroids on the metabolism of calcium, phosphorus, and vitamin D (44). In such cases, the routine diet should contain sufficient calcium and phosphorus (1200 mg/day), and their serum levels should be monitored regularly. Furthermore, phosphorus supplementation may be required to correct hypophosphatemia (45).

Hydration status should also be closely monitored after kidney transplantation; since most kidney transplant patients require fluid restriction during dialysis, they should be reminded by a nutritionist to receive adequate fluids after transplantation (Table 2) (45, 46). Most of the patients who receive a kidney transplant have high triglycerides or cholesterol due to multifactorial issues, and interventions such as calorie restriction are recommended for overweight or obese patients. Among the other strategies in this regard are cholesterol intake of less than 300 mg/day and restricted total fat intake (47).

Conclusion

Nutrition therapy and counseling play a key role in the recovery after transplantation in patients with renal failure. It seems that nutritional and medical evaluation based on laboratory methods, vital signs, and anthropometric measurements are essential to improving the quality of nutritional interventions after kidney transplantation, while these measures also contribute to reducing the risk of kidney transplant rejection. Therefore, the periodic evaluation and follow-up of these patients may yield beneficial outcomes.

Authors' Contributions

S. H. Z., N. P., F. Y., and A. N. designed the research; S. H. Z., N. P., and A. N. conducted the library search and drafted the manuscript; N. P. and F. Y. designed the tables; A. N., S. H. Z., and N. P. participated in the drafting and editing of the manuscript. All the authors read and approved the final manuscript.

Acknowledgments

Hereby, we extend our gratitude to our colleagues for assisting us in the nutritional assessments and kidney transplantation procedures for their valuable comments.

Funding/Support

None

Conflicts of Interest

None declared

References

- McDonald SP, Russ GR. Survival of recipients of cadaveric kidney transplants compared with those receiving dialysis treatment in Australia and New Zealand, 1991–2001. *Nephrology Dialysis Transplantation*. 2002;17(12):2212-9.
- Zhang Q-L, Rothenbacher D. Prevalence of chronic kidney disease in population-based studies: systematic review. *BMC Public Health*. 2008;8(1):117.
- Chan M, Patwardhan A, Ryan C, Trevillian P, Chadban S, Westgarth F, et al. Evidence-based guidelines for the nutritional management of adult kidney transplant recipients. *J Ren Nutr*. 2011;21(1):47-51.
- Veroux M, Corona D, Sinagra N, Tallarita T, Ekser B, Giaquinta A, et al. Nutrition in kidney transplantation. *Int J Artif Organs*. 2013;36(10):677-86.
- Tritt L. Nutritional assessment and support of kidney transplant recipients. *J Infus Nurs*. 2004;27(1):45-51.
- Teger NB. Owner's Manual: Nutrition Care for Your Kidney Transplant. *J Ren Nutr*. 2019;29(3):249-55.
- Ebrahimi F, Aryaeian N, Pahlavani N, Abbasi D, Hosseini AF, Fallah S, et al. The effect of saffron (*Crocus sativus* L.) supplementation on blood pressure, and renal and liver function in patients with type 2 diabetes mellitus: A double-blinded, randomized clinical trial. *Avicenna J Phytomed*. 2019;9(4):322.
- Nieto T, Inston N, Cockwell P. Renal transplantation in adults. *Bmj*. 2016;355:i6158.
- Zrim S, Furlong T, Grace BS, Meade A. Body mass index and postoperative complications in kidney transplant recipients. *Nephrology*. 2012;17(6):582-7.
- Cederholm T, Bosaeus I, Barazzoni R, Bauer J, Van Gossum A, Klek S, et al. Diagnostic criteria for malnutrition—an ESPEN consensus statement. *Clin Nutr*. 2015;34(3):335-40.
- Hasse JM. Nutrition assessment and support of organ transplant recipients. *J Parenter Enteral Nutr*. 2001;25(3):120-31.
- Kaya E, Bakir A, Koseoglu YK, Velidedeoglu M, Trabulus S, Seyahi N. Association of Nutritional Assessment by Phase Angle With Mortality in Kidney Transplant Patients in an 8-Year Follow-Up. *Progress in Transplantation*. 2019;29(4):321-6.
- Gibson RS. *Principles of nutritional assessment*: Oxford university press, USA; 2005.
- Javid Z, Shadnoush M, Khadem-Rezaiyan M, Honarvar NMZ, Sedaghat A, Hashemian SM, et al. Nutritional adequacy in critically ill patients: Result of PNSI study. *Clin Nutr*. 2020.
- Silva MIB, da Silva Lemos CC, Torres MRSG, Bregman R. Waist-to-height ratio: an accurate anthropometric index of abdominal adiposity and a predictor of high HOMA-IR values in nondialyzed chronic kidney disease patients. *Nutrition*. 2014;30(3):279-85.
- Group WW. Use and interpretation of anthropometric indicators of nutritional status. *Bulletin of the World health organization*. 1986;64(6):929.

17. Ness-Abramof R, Apovian CM. Waist circumference measurement in clinical practice. *Nutr Clin Pract.* 2008;23(4):397-404.
18. Zhang J, Cao X, Ping S, Wang K, Shi J, Zhang C, et al. Comparisons of ethanol extracts of Chinese propolis (poplar type) and poplar gums based on the antioxidant activities and molecular mechanism. *J Evid Based Complementary Altern Med.* 2015;2015.
19. Lee Y, Shin D-h, Kim J-H, Hong S, Choi D, Kim Y-J, et al. Caffeic acid phenethyl ester-mediated Nrf2 activation and IκB kinase inhibition are involved in NFκB inhibitory effect: structural analysis for NFκB inhibition. *Eur J Pharmacol.* 2010;643(1):21-8.
20. Chan A-W, Tetzlaff JM, Altman DG, Laupacis A, Gøtzsche PC, Krleža-Jerić K, et al. SPIRIT 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med.* 2013;158(3):200-7.
21. Excler JL, Sann L, Lasne Y, Picard J. Anthropometric assessment of nutritional status in newborn infants. Discriminative value of mid arm circumference and of skinfold thickness. *Early Hum Dev.* 1985;11(2):169-78.
22. Buffa R, Mereu E, Succa V, Latini V, Marini E. Specific BIVA recognizes variation of body mass and body composition: Two related but different facets of nutritional status. *Nutrition.* 2017;35:1-5.
23. Roudi F, Khosravi M, Pahlavani N. A narrative review of general principles of nutrition status assessment in hospitalized adult patients. *Navid No.* 2019;21(68):62-76.
24. Affret A, Wagner S, El Fatouhi D, Dow C, Correia E, Niravong M, et al. Validity and reproducibility of a short food frequency questionnaire among patients with chronic kidney disease. *BMC nephrology.* 2017;18(1):297.
25. Bross R, Noori N, Kovesdy CP, Murali SB, Benner D, Block G, et al., editors. *Dietary assessment of individuals with chronic kidney disease. Seminars in dialysis;* 2010: Wiley Online Library.
26. Mekki K, Remaoun M, Belleville J, Bouchenak M. Hemodialysis duration impairs food intake and nutritional parameters in chronic kidney disease patients. *Int Urol Nephrol.* 2012;44(1):237-44.
27. Desbrow B, Bauer J, Blum C, Kandasamy A, McDonald A, Montgomery K. Assessment of nutritional status in hemodialysis patients using patient-generated subjective global assessment. *J Ren Nutr.* 2005;15(2):211-6.
28. Steiber AL, Kalantar-Zadeh K, Secker D, McCarthy M, Sehgal A, McCann L. Subjective Global Assessment in chronic kidney disease: a review. *J Ren Nutr.* 2004;14(4):191-200.
29. Pahlavani N, Sadeghi A, Rasad H, Azizi Soleiman F. Relation of inflammation and oxidative stress with blood glucose, lipids and BMI, fat mass and body weight in people with type 2 diabetes. *Diabetes Nurs.* 2014;2(2):42-51.
30. Sauberlich HE. *Laboratory tests for the assessment of nutritional status:* Routledge; 2018.
31. Maeda H, Sogawa K, Sakaguchi K, Abe S, Sagizaka W, Mochizuki S, et al. Urinary albumin and transferrin as early diagnostic markers of chronic kidney disease. *J Vet Med Sci.* 2015:14-0427.
32. Kamal A. Estimation of blood urea (BUN) and serum creatinine level in patients of renal disorder. *Indian J Fundam Appl Life Sci.* 2014;4(4):199-202.
33. Jansz TT, Bonenkamp AA, Boereboom FTJ, van Reekum FE, Verhaar MC, van Jaarsveld BC. Health-related quality of life compared between kidney transplantation and nocturnal hemodialysis. *PLoS One.* 2018;13(9):e0204405-e.
34. Mahan LK, Escott-Stump S, Krause MV. *Krause's food & nutrition therapy:* Elsevier Saunders; 2007.
35. Andrews LM, Li Y, De Winter BC, Shi Y-Y, Baan CC, Van Gelder T, et al. Pharmacokinetic considerations related to therapeutic drug monitoring of tacrolimus in kidney transplant patients. *Expert Opinion on Drug Metabolism & Toxicology.* 2017;13(12):1225-36.
36. Nikkel L, Mohan S, Zhang A, McMahon D, Boutroy S, Dube G, et al. Reduced fracture risk with early corticosteroid withdrawal after kidney transplant. *Am J Transplant.* 2012;12(3):649-59.
37. Ciriaco M, Ventrice P, Russo G, Scicchitano M, Mazzitello G, Scicchitano F, et al. Corticosteroid-related central nervous system side effects. *Journal of pharmacology & pharmacotherapeutics.* 2013;4(Suppl1):S94.
38. Pahlavani N, Roudi F, Zakerian M, Ferns GA, Navashenaq JG, Mashkouri A, et al. Possible molecular mechanisms of glucose-lowering activities of *Momordica charantia* (karela) in diabetes. *J. Cell. Biochem.* 2019;120(7):10921-9.
39. Teplan V, Valkovsky I, Teplan Jr V, Stolova M, Vyhnanek F, Andel M. Nutritional consequences of renal transplantation. *J Ren Nutr.* 2009;19(1):95-100.
40. Pahlavani N, Malekhamdi M, Firouzi S, Rostami D, Sedaghat A, Moghaddam AB, et al. Molecular and cellular mechanisms of the effects of Propolis in inflammation, oxidative stress and glycemic control in chronic diseases. *Nutrition & Metabolism.* 2020;17(1):1-12.
41. Pahlavani N, Sedaghat A, Moghaddam AB, Kiapey SSM, Navashenaq JG, Jarahi L, et al. Effects of propolis and melatonin on oxidative stress, inflammation, and clinical status in patients with primary sepsis: Study protocol and review on previous studies. *Clin Nutr ESPEN.* 2019;33:125-31.
42. Moore MC. *Mosby's Pocket Guide to Nutritional Assessment and Care-E-Book:* Elsevier Health Sciences; 2016.
43. Nolte Fong JV, Moore LW. Nutrition trends in kidney transplant recipients: the importance of dietary monitoring and need for evidence-based recommendations. *Frontiers in medicine.* 2018;5:302.
44. Dizdar OS, Yıldız A, Gul CB, Gunal AI, Ersoy A, Gundogan K. The effect of hemodialysis, peritoneal dialysis and renal transplantation on nutritional status and serum micronutrient levels in patients with end-stage renal disease; multicenter, 6-month period,

longitudinal study. *J Trace Elem Med Biol.* 2020;126498.

45. Soldavini J. Krause's Food & The Nutrition Care Process. *J Nutr Educ Behav.* 2019;51(10):1225.

46. Kruszyna T, Niekowal B, Kraśnicka M, Sadowski J, editors. Enhanced recovery after kidney

transplantation surgery. *Transplantation proceedings;* 2016: Elsevier.

47. Loncar D, Tulumovic E, Bijedic I, Bijedic A. Hyperlipidemia and cardiovascular diseases in patients with transplanted kidney. *Atherosclerosis.* 2018;275:e216.